

AN ESSAY  
ON  
ANIMAL, MINERAL, AND VEGETABLE  
POISONS.

WITH THIRTEEN COLOURED PLATES.


*Price 3s. 6d. bds.*

21857 / A

C. XVII  
10/

ESSAY (AM)  
C

41022




AN  
ESSAY

ON

*Mineral, Animal, and Vegetable*

POISONS.







*Digitalis*  
*Purpurea*

*Foxglove*



AN  
**ESSAY**  
ON  
**MINERAL, ANIMAL, AND VEGETABLE**  
**POISONS ;**

IN WHICH  
THE SYMPTOMS, MODE OF TREATMENT AND  
TESTS OF EACH PARTICULAR POISON,  
WITH THE  
GENERAL MORBID APPEARANCES *on* DISSECTION,  
ARE CONCISELY DETAILED :  
TO WHICH IS ADDED,  
AN ACCOUNT OF THE MEANS TO BE EMPLOYED  
IN CASES OF SUSPENDED ANIMATION.

**Third Edition,**

*Illustrated by Thirteen Coloured Plates.*

---

PRINTED FOR

*Cox and Son, St. Thomas's Street, Southwark ;*

Also by T. and G. Underwood, and Highley, Fleet Street ;  
Callow and Wilson, Prince's Street, Soho ; Simpkin and  
Marshall, Stationers' Court ; Anderson, West  
Smithfield ; John Cox, Berners Street,  
Oxford Street ; and W. Stewart,  
Edinburgh.

---

1823.



TO JAMES BLUNDELL, M. D.

LECTURER

ON

PHYSIOLOGY AND MIDWIFERY

AT

GUY'S HOSPITAL.

SIR,

THIS little work having originally appeared under your auspices, I feel ambitious to solicit the same patronage for the present edition, which indeed differs but slightly from the former.

It is gratifying to my feelings that an opportunity is thus afforded of expressing the high opinion I entertain of your public and private character.

That you may long continue to enjoy increasing degrees of professional celebrity, as the just reward of unwearied diligence and perseverance,  
is the wish of

Sir,

Your most obedient and obliged servant,  
THE AUTHOR.

LONDON.



## INTRODUCTION.

---

SEVERAL reasons have induced the author of the following pages to trespass on the attention of the public ; more particularly, however, the importance of the subject under consideration, and the repeated and pressing solicitations of his friends. These apologies would ill suffice, if a small work on the subject of Poisons were already in print ; but there being none expressly confined to this subject, in a concise form, was another strong inducement for this undertaking. Conciseness, as far as the nature of the subject would allow, without perplexity, has been particularly studied. In addition to the remarks

on poisons, the author has subjoined a few observations on the means to be had recourse to in cases of suspended animation, knowing this to be an accident of not unfrequent occurrence, and one requiring immediate assistance. He is now willing to offer it to the inspection of the public ; and if it should be the means of saving one of his fellow creatures from an untimely end, his trouble will be fully recompensed.

# ON POISONS.



**D**IFFERENT plans have been adopted by authors in the arrangement of Poisons, each following some system peculiar to himself; that pursued by Orfila is the most scientific; but we shall, in this little work, deviate from his system, and class them according to the three kingdoms:—Mineral, Vegetable, and Animal.



## MINERAL POISONS.



### *CORROSIVE METALLIC SALTS.*

**T**HE symptoms which follow an over-dose of the more corrosive metals, are very similar in the different metallic salts. The urgency of the

symptoms will necessarily depend on the quantity taken—the form, whether solid or fluid—the state of the stomach at the time, and other occasional circumstances. We shall relate the general operations of this class of poisons, and then the particular poisons, and their peculiarities.

When a person has taken a sufficient quantity of any substance to produce deleterious effects upon the constitution, it is said to be an over-dose, or in other words to act as a poison.

### *General Symptoms.*

If Arsenic or Corrosive Sublimate be swallowed, it occasions sickness and uneasiness about the stomach, violent retchings, sense of heat about the mouth and fauces, with a disagreeable taste; the pain in the stomach then becomes very distressing, and blood is sometimes ejected; the bowels soon become affected, and a discharge of offensive matter takes place, frequently mixed with blood, and accompanied with considerable griping and



tenesmus. The countenance becomes anxious ; the breathing difficult ; thirst excessive ; skin hot ; and pain at the stomach and bowels much increased, particularly upon pressure : then cold sweats alternating with flushes of heat, cold extremities, faintness, convulsions, and the most distressing symptoms ; which are soon followed by death, relieving the patient from an exquisite state of misery. The pulse is usually small, quick, and irregular, but at other times is scarcely affected ; and is therefore not to be depended on.

---

## ARSENICAL PREPARATIONS.

---

*OXYD of ARSENIC, or ARSENIUS ACID : (Arsenici Oxydum.)*

THIS mineral, and all the preparations obtained from it, are highly poisonous, even in very small quantities ; from which circumstance, it is of the greatest consequence its effects should

be watched, when administered for the cure of any disease; and whenever any distressing symptoms make their appearance, it should be immediately discontinued. The same observation will apply to all poisons, particularly the more active minerals; i. e. when any unpleasant symptoms, peculiar to the poison at that time employed, show themselves, it should be suspended for a time.

The Oxyds and Sulphurets of Arsenic, and Arseniates, possess poisonous qualities in different degrees, and will all destroy life if the dose be at all considerable. Fly Water is commonly a Solution of this mineral.

### *Symptoms.*

For these we will refer the reader to the head of *Corrosive Metallic Salts*. They are generally very distressing, and it is a poison, the operation of which is very energetic in the majority of instances; cases have, however, occurred, where patients have been destroyed without the production of any distressing symptoms,

where very large quantities have been taken. One symptom peculiar to this poison is a copious flow of saliva, not having the mercurial fœtur : the evacuations are often green.

### *Treatment.*

Unless our attention is directed to the patient early, there will be little chance of success. As we possess no antidote to this poison, our object is to expel it immediately from the stomach by emetics, should it not have excited vomiting, which it scarcely ever fails of doing. Milk, white of eggs, and mucilaginous drinks should be taken freely, to encourage the vomiting and cleanse out the stomach. Emollient Clysters are also to be given, to remove any of the Arsenic which may have escaped into the intestines. It is useless losing time in administering chemical salts, under the idea of neutralizing the effects of the poisons; we have no substance possessed of that power. Sulphurets of Potass and Soda, Lime Water, and Alkalies, have all been used without success. **Inflammatory**

symptoms must be combated by the usual anti-phlogistic means.

The *external application* of Arsenic will give rise to all the above symptoms, and destroy life.

### *Tests.*

1. A solution of Arsenic is changed yellow by the addition of water saturated with sulphuretted Hydrogen.

2. With Sulphate of Copper and Caustic Potass or Ammonia it forms a beautiful green precipitate of *Arseniate of Copper*.

3. By adding to it a small quantity of Liquor Ammonia and a Solution of Nitrate of Silver, you will produce a beautiful yellow precipitate of *Arseniate of Silver*.

4. The above results will not, however, enable you to speak decisively as to the presence of this mineral. It is necessary to reduce some to its metallic state, before we have clear evidence of its existence in any fluid. If any should be rejected solid from the stomach, mix it with some Potass and Charcoal, then submit the mixture to

the heat of a candle or spirit lamp, in a glass tube, and the Metallic Arsenic will be sublimed, and condense itself on the upper part of the tube, in cubic crystals. This is a property possessed by no other metal. If there should not be any powder in the matter vomited, or in the contents of the stomach (supposing the patient dead), the precipitate obtained in Test 2, or 3, may be submitted to the same process, and a similar result will follow.

---

## MERCURIAL PREPARATIONS.

---

### *CORROSIVE SUBLIMATE:*

*(Hydrargyri Oxymurias.)*

#### *Symptoms.*

IN addition to the general symptoms enumerated above, we may notice that the pain of the stomach and abdomen, is generally violent in the extreme; the bowels are speedily deranged, and the evacuations bloody: salivation, with the

peculiar mercurial factor, is another well-marked symptom, and one early in its appearance ; for it may be remarked here, that none of the preparations of Mercury affect the salivary glands in so short a time as this.

### *Treatment.*

White of eggs and milk, immediately ; the former decomposes Corrosive Sublimate, and throws down an insoluble salt, comparatively mild in its operation. Barley water and linseed tea, or any other mucilaginous fluid, to be taken freely to wash out the stomach effectually. The bowels are next to be attended to ; give saline purgatives by the mouth, with emollient clysters. The inflammation must not be overlooked, but treated by leeches, blisters, &c. ; and, when the patient rallies somewhat, by venesection. General bleeding will not be proper immediately on the appearance of the inflammation in many cases.

### *Tests.*

1. With Corrosive Sublimate, Albumen

forms a white precipitate of *Submuriate of Mercury*; this test will detect very minute quantities of the Salt.

2. Alkalies form with it a red or yellowish precipitate.

3. *Liquor Ammoniaë* gives a white precipitate.

4. Reduction of the metal with flux (*Potass and Charcoal*), is the only decisive test.

Some other preparations of Mercury act in a similar manner, but with much less violence; such as the Red Oxyd, Nitrates, Sulphurets, &c. The plan of treatment would be similar, and the tests the reduction of the metal.

---

## ANTIMONIAL PREPARATIONS.

---

*EMETIC TARTAR: (Antimonium  
Tartarizatum.)*

*Symptoms.*

THE vomiting, which is the first symptom,

is very distressing and urgent; it is soon followed by pain in the stomach; spasm of the œsophagus; great prostration of strength; a quick small pulse: colicky pains soon attack the bowels, and violent purging succeeds; breathing, anxious and hurried; cold perspirations and other symptoms as above.

### *Treatment.*

Decoction of Oak Bark, Infusion of Galls, and common Tea, all decompose the Tartar Emetic; and should be taken freely, that it may be thoroughly expelled from the stomach; mucilaginous drink and milk are also proper. If the stomach should continue very irritable, give Opiates. — Treat other symptoms as usual.

### *Tests.*

1. With Sulphuric Acid, Lime Water, and the Alkalies, it forms a white precipitate.

2. A Decoction of Oak Bark or Galls, throws down a yellowish precipitate.



**BUTTER OF ANTIMONY:***(Antimonii Murias.)**Symptoms.*

This acts as a powerful escharotic, and when swallowed causes great destruction, giving rise to symptoms of a very distressing nature, as violent as are observed after the Corrosive Sublimate has been taken.

*Treatment.*

Same as for Tartar Emetic.

*Tests.*

1. The revival of the metal, by flux.
2. It forms a white flaky precipitate with water.

The Antimonial Wine, which is often administered by nurses to children, has not unfrequently proved a destructive poison, killing them insidiously; this, the Sulphurets, and all Antimonial Preparations, act in a similar manner, and the symptoms they occasion must be treated as for the Tartar Emetic.

## PREPARATIONS OF COPPER.

---

### *VERDIGRIS : (Subacetas Crupri.)*

THE preparations of copper are seldom taken or given intentionally as poisons ; but from neglect and want of cleanliness, in leaving acid and vegetable substances in copper vessels, these sometimes become corroded, and the Oxyd or Carbonate of Copper is mixed with the food.

#### *Symptoms.*

The Salts of Copper give rise to colicky pain in the stomach and bowels ; nausea ; coppery taste ; vomiting greenish matter, and violent head-ache ; severe griping pain in the bowels, with purging often of blood ; these symptoms are sometimes succeeded by cold perspirations, convulsions, and death.

#### *Treatment.*

Milk, white of eggs, sugared water, and mu-

milaginous fluids freely. If the spasms of the alimentary canal be severe, Opiates will be useful; Emollient Clysters to evacuate and lubricate the bowels.

The Sulphate, Nitrate, Muriate, Carbonate and Oxyds, of Copper, and Wines in which either of these preparations may be present, act in a similar way and require the same treatment.

### *Tests.*

1. All the Salts of Copper are of a green or blue colour.

2. Liq. Ammoniæ, when added to them, forms a greenish precipitate; but if added in excess, the precipitate becomes dissolved, and a beautiful blue solution of the *Ammoniaret of Copper* is the result. If the salt be so much diluted as not to colour the water, this test will detect it.

3. Prussiate of Potass forms a brown precipitate.

## PREPARATIONS OF TIN.

---

*MURIATE OF TIN: ) Stanni Murias. )*

### *Symptoms.*

VIOLENT vomiting and purging, with spasmodic affections of the stomach and bowels; cramps; sharp quick pulse; sometimes paralysis; with convulsions, and death. It has been mistaken for Epsom Salts, and caused death.

### *Treatment.*

Emetic, if necessary; afterwards milk and mucilaginous fluids; Emollient and Oily Clysters. Opiates to allay spasm of the bowels, &c.: antiphlogistic treatment, if required.

### *Tests.*

1. With Nitro-Muriate of Gold, it forms a brownish red, or purplish precipitate, (the *Powder of Cassius.*)

2. With Prussiate of Potass, a white precipitate.

All the preparations of Tin resemble this in their effects, &c.



## PREPARATIONS OF ZINC.

---

*WHITE VITRIOL: (Zinci Sulphas.)*

### *Symptoms.*

It is generally rejected, immediately that it reaches the Stomach, hence it rarely destroys life. It produces a peculiar astringent metallic taste, with a sense of suffocation; but vomiting soon relieves these symptoms; otherwise the countenance becomes pallid and sunk; pulse quick and irregular; bowels always more or less affected: sometimes symptoms occur resembling the Lead Colic, which are succeeded by obstinate diarrhœa.

*Treatment.*

Alkalies, Magnesia, or Chalk, are the best remedies : After this diluents and clysters, with Opium if much spasmodic colic.

*Tests.*

1. Forms with Alkalies a white precipitate, which is readily dissolved by Sulphuric Acid.

2. With Prussiate of Potass, a blue precipitate.

3. With Chromate of Potass, an orange yellow precipitate.



## PREPARATIONS OF SILVER.

---

### *LUNAR CAUSTIC : ( Argenti Nitras. )*

*Symptoms.*

It is one of the most corrosive poisons ; as we might expect from the result of its external ap-

plication. It is, fortunately, rarely taken as a poison. Its action is similar to the *Corrosive Sublimate*.

### *Treatment.*

The best antidote is mucilaginous fluids, in which Common Salt has been dissolved; this forms a Muriate of Silver, insoluble and harmless: in other respects treat it as for *Corrosive Sublimate*.

### *Tests.*

1. With Alkalies, it forms a white precipitate.
2. With Muriatic Acid, and Saline Muricates, a white precipitate, which soon changes to a blackish colour, by exposure to the air; it may be dissolved by Ammonia.

## GENERAL MORBID APPEARANCES.

When life is destroyed by the metallic salts above enumerated, dissection proves that the appearances are similar in the majority of cases. Where *Arsenic* has killed, the stomach and

intestines are highly inflamed, often with gangrenous spots in different parts of the former viscus; the villous coat sometimes becomes eroded, and so tender as to be readily peeled off; the peritoneum, throughout the abdomen, is frequently in a state of inflammation. If death has resulted from *Corrosive Sublimate*, the inflammation of the villous surface of the stomach and intestines is more general, and the intestines are sometimes ulcerated. In death from *Tartar Emetic*, the morbid appearances are not generally sufficient to account for dissolution. In the other Salts which have caused death, the appearances have resembled those from *Corrosive Sublimate*; but there are no diagnostic marks by which we can tell what poison has destroyed our patient: it may therefore be sufficient to say, that all the Metallic Salts mentioned above, produce more or less inflammation of the stomach and intestines, and of the other abdominal viscera.



## PREPARATIONS OF LEAD.

---

### *SUGAR OF LEAD : (Plumbi Superacetas)*

#### *Symptoms.*

THIS metal produces effects upon the constitution in a great measure peculiar to itself, giving rise to considerable derangement in the nervous system ; as is frequently observed in plumbers and painters. In a large dose it occasions pain at the stomach ; an astringent metallic taste ; sometimes vomiting ; obstinate constipation ; colicky pains in the bowels ; and contractions of the abdominal muscles : these are succeeded by pallid countenance ; tremors ; sometimes delirium ; and if the patient should survive the primary symptoms, paralytic affections seldom fail to make their appearance.

#### *Treatment.*

Emetics ; Sulphate of Magnesia, dissolved in

mucilaginous fluids ; Opiates, to allay the spasm of the bowels ; warm bath ; Castor Oil, and purgative clysters, composed of Infusion of Colocynth, or Senna and Salts.

### *Tests.*

1. Sulphuric Acid, and the Alkaline Sulphate, form a white insoluble precipitate.

2. Chromate of Potass, throws down a yellow precipitate.

3. Sulphuretted Hydrogen, forms at first a brown precipitate, but it soon becomes black.

4. Carbonated Alkalies, form a white precipitate.

### *Morbid Appearances.*

There is merely a stricture about the Colon, or rather a general contraction of that intestine ; no marks of inflammation are observed.

The Subacetate, Carbonate, and Oxyds of Lead, and Wines or other fluids, which are either accidentally or purposely impregnated with preparations of Lead, give rise to symptoms

similar to those from *Sugar of Lead*, and require the same treatment.

The other Metallic Salts, as those of Gold, Platina, Bismuth, Nickel, &c. are never employed as poisons.

---

## CAUSTIC ALKALIES, AND THEIR SUBCARBONATES.

---

### *POTASS, SODA, AND AMMONIA.*

#### *Symptoms.*

DISTRESSING heat and pain in the fauces, œsophagus, and stomach ; nausea ; an urinous caustic taste ; vomiting, often of blood ; intestines soon become affected, and bloody evacuations are the result. *Ammonia* is the most active in its operation, generally causing convulsions and derangement of the intellectual faculties. A

large dose of Liquid Ammonia has destroyed life in five minutes.

### *Treatment.*

Similar in all cases ; neutralize the Alkali by some mild acid, as vinegar ; diluted lemon juice, &c. and give mucilaginous drinks.

### *Tests.*

Alkalies have an urinous taste ; they change violets green, and turmeric paper brown ; precipitate Metallic Oxyds, from their solutions ; and form soapy compounds with oily substances.

1. *Ammonia* has a pungent odour.
2. It changes the salts of copper blue.

*Potass and Soda* may be distinguished by the former becoming precipitated in a state of *Cream of Tartar*, when Tartaric Acid is added in excess ; whilst soda forms a very soluble compound, with Tartaric Acid in any quantity.

A solution of Platina throws down a yellow precipitate with *Potass* and not with *Soda*.

*Morbid Appearances.*

Inflammation of the œsophagus, stomach, and bowels, which frequently present a gangrenous and sloughy appearance.

---

## SALTS OF BARYTES.

---

### MURIATE AND NITRATE OF BARYTES.

*Symptoms.*

VOMITING; purging; violent pain in the stomach and bowels; vertigo; insensibility, paralysis; convulsions; and death. They act with considerable violence and activity.

*Treatment.*

Sulphate of Soda, of Magnesia, or of Potass, dissolved in some mucilaginous fluid, to be drank freely: they form an insoluble Sulphate, which is inert.

*Tests.*

The Salts of Barytes form an insoluble compound with Sulphuric Acid and its salts, which will detect very minute quantities of this earth.

*Morbid Appearances.*

Same as from the Alkalies.

The Carbonate, Acetate, and other soluble Salts of Barytes, act in a similar manner, and require precisely the same treatment.

---

## MINERAL ACIDS.

---

*OIL of VITRIOL: (Acidum Sulphuricum.)*

*AQUA FORTIS: (Acidum Nitricum )*

*SPIRITS of SALTS: (Acidum Muriaticum.)*

*Symptoms.*

VIOLENT burning pain about the throat, fauces, œsophagus, and whole alimentary canal; fœtid eructations; vomiting of blood; pulse

very small and irregular ; abdomen tense ; great thirst, cold perspirations ; convulsions, and death.

### *Treatment.*

In all cases where either of the acids above mentioned have been swallowed, our antidotes must be immediately administered, or they will be useless. *Calcined Magnesia* is the best remedy : if not at hand, soap, chalk, or the alkalies diluted.

### *Tests.*

They turn vegetable blues red. Form neutral Salts with the Alkalies, and effervesce when added to a carbonated alkali or earth.

*Sulphuric Acid*, forms a white insoluble precipitate, both with Muriate of Barytes and the Acetate of Lead.

*Muriatic Acid*, forms an insoluble precipitate with *Nitrate of Silver*. If the fumes arising from it come in contact with Ammoniacal gas, a dense white vapour of *Muriate of Ammonia*, is the result.

*Nitric Acid* may be known by its red fuming nature ; by its forming Nitre when saturated with Potass ; and if paper be dipped into the saturated solution, it is converted into touch-paper. It forms no precipitate with the salts used to test the other acids.

### *Morbid Appearances.*

If the Acids be concentrated, they occasion complete disorganization and destruction of the parts with which they come in contact: the lining membrane of the œsophagus, stomach, and intestines, is in a high state of inflammation, and holes are generally formed in the stomach, with gangrenous appearances surrounding them. The *Nitric Acid* proves most destructive; the stomach is converted into a pulpy, soft, blackish, ragged substance, and completely altered in its character ; and from the escape of the Acid, the peritoneum becomes inflamed ; and where this acid has killed, the parts which it has come in contact with, are yellow.

The *Oxalic*, *Tartaric*, *Nitric*, and *Fluoric*



*Acids*, when taken in sufficient quantity and undiluted, will give rise to very violent symptoms, and even destroy life; several instances of which have unfortunately happened by the *Oxalic Acid*, or *Acid of Sugar*.

### *Tests.*

*Oxalic Acid* very much resembles Epsom Salts in its external appearance. It is excessively sour to the taste. It readily detects very minute quantities of the Salts of Lime, forming a white insoluble precipitate with them. The crystals are four-sided prisms.

*Tartaric Acid* forms a compound of Supertartrate of Potass, when added in excess to a solution of Potass.

*Citric Acid* crystallizes in rhomboidal prisms, and is a basis of Lemon Juice.

*Fluoric Acid* has a suffocating smell, and possesses the property of corroding glass, acting upon the silex which it contains.

These, with some other acids which it will be

unnecessary to enumerate, produce symptoms, &c., similar to the Mineral Acids, and require a similar mode of treatment, viz. Magnesia, Chalk, &c.

*SALT PETRE : (Potassæ Nitras.)*

*Symptoms.*

Violent pain in the stomach, with spasm ; vomiting and purging of blood : very irregular pulse ; great prostration of strength ; syncope ; coldness of the extremities ; clammy perspiration ; involuntary stools, and speedy dissolution : if the primary symptoms should not destroy, the nervous system becomes more particularly deranged, and paralysis is the consequence.

*Treatment.*

Emetics, mucilaginous drinks, milk, and emollient purgatives and clysters.

*Tests.*

1. Crystals six-sided prisms.
2. Detonates on burning coals.

3. When mixed with Sulphuric Acid, red nitrous fumes escape.

### *Morbid Appearances.*

Same as are observed in cases of death from the Corrosive Metals.

## *PHOSPHORUS.*

### *Symptoms.*

Most distressing pain and heat in the stomach, which are more urgent if the poison be dissolved; besides this, it occasions other symptoms as observed in the Corrosive Metals.

### *Treatment.*

It is advisable to expel it from the stomach as speedily as possible; it has been recommended to distend the stomach with food, and then to excite vomiting and give diluents freely. The operation is less violent, the more it is excluded from the air.

*Tests.*

Its peculiar odour, and general properties, will commonly enable any person to detect it. If dissolved in oil, it is luminous in a dark room.

*Morbid Appearances.*—See Corrosive Metals.

---

## VEGETABLE POISONS.

---

### NARCOTICS.

#### OPIUM AND ITS PREPARATIONS

##### *Symptoms.*

THIS is one of the most common and destructive poisons of this class, and produces symptoms common to all vegetable narcotics. A full dose of *Opium* occasions almost immediate insensibility; with slow pulse; stertorous breathing; dilated pupil: greatest difficulty in being roused: the countenance is at

first flushed, but soon becomes pallid; sometimes convulsions and paralysis precede dissolution. Vomiting is not an usual symptom when the dose is large.

### *Treatment.*

Sulphate of Zinc, gr. viij. vel gr. x. every ten minutes, till vomiting is excited: or Sulphate of Copper, gr. j. to gr. iij.; tickle the throat with a feather, and use every possible means to evacuate the stomach: it will be advisable to pass an elastic tube down the œsophagus and thus introduce emetics, if the patient cannot swallow: the person must be kept in constant motion and exercise by able assistants. In some cases it will be proper to open the jugular vein, to relieve the vessels of the brain from a state of congestion. If scarcely any pulse, Wine, Brandy, and Ammonia, should be introduced into the stomach. Acids should never be given till we are thoroughly convinced no Opium remains in the stomach. Active purgatives both by the mouth and per anum. Coffee may be

drank freely. Saline purgatives and effervescing draughts are useful when the patient is recovering.

Children are often very much injured, and sometimes destroyed, by the too free use of *Syrup of Popies*; it give rise to drowsiness, insensibility, and convulsions. Treatment here will be Ammonia. Brandy diluted, and Wine with the warm bath.

We are possessed of no tests by which we can distinguish poisons of this class, and can only conjecture they have been taken, by their taste or smell, and the symptoms.

In general no morbid appearances are evident.

HENBANE (*Hyoscyamus Niger*,) HEMLOCK (*Conium Maculatum*) STRONG-SCENTED LETTUCE (*Lactuca Virosa*,) WATER HEMLOCK (*Cicuta Virosa*,) DEADLY NIGHTSHADE (*Atropa Belladonna*.)

These, with some others, produce symptoms very similar to Opium, and require the same mode of treatment.

MONKSHOOD (*Aconitum Napellus*,) POISON NUT (*Nitx Vomica*) SPURIOUS ANGUSTURA BARK (*Angustura Pseudo Ferruginea*,) CAMPHOR (*Laurus Camphora*,) POISONOUS FUNGI, with some other, both indigenous and exotic poisons, produce symptoms in some measure resembling those occasioned by the simple Narcotics; they however excite more acrimony and disturbance in the alimentary canal; some exhilarating effects precede their sedative operation, unless the dose be very large; and they more commonly occasion convulsions and spasmodic affections. These have been arranged, by Orfila, under the class Acro-Narcotics.

### *Treatment.*

As for Narcotics. If the breathing be suspended, keep up artificial respiration.

### *ACRID VEGETABLES.*

BLACK AND STINKING HELLEBORE (*Helleborus Niger et Fætidus*) WHITE HELLEBORE (*Veratrum Album*), SQUIRTING

CUCUMBER (*Momordica Elaterium*), GAMBOGE (*Stalagmitis Cambogioides*), EUPHORBIA, *several species*; SAVINE (*Juniperus (Sabina)*), MEADOW SAFFRON (*Colchicum Autumnal*), SQUILL (*Scilla Maritima*), *several species* of RANUNCULUS, &c. &c.

### *Symptoms.*

All these vegetables, in over-doses, and many others, occasion an acrid taste ; burning sensation about the mouth and fauces ; constriction of the fauces ; pain in the stomach and bowels ; distressing vomiting and purging, often of blood, succeeded by insensibility, difficulty and shortness of breath, and convulsions.

### *Treatment.*

Mild emetics, mucilaginous diluents, milk, emollient laxatives, and clysters. If the vomiting, after the poison has been thoroughly expelled, should continue to distress the patient, mild opiates are proper.



*Morbid Appearances.*

Sometimes, in the more acrid vegetables, there is some inflammation of the stomach and bowels ; but we cannot generally depend upon these appearances in poisoning from vegetable substances.

*PRUSSIC ACID.**Symptoms.*

It occasions nausea, vertigo, vomiting, sense of suffocation, and almost immediate death, if concentrated. The mere act of smelling to it produces a sense of suffocation and giddiness. Fortunately this poison is seldom taken in the concentrated form ; but from its existence in the kernels of some fruits, laurel leaves, &c. it is right to be acquainted with its operation. In one case it caused immediate death, as if the patient had been suddenly seized with an apoplectic fit. Its external application will destroy life. The essential oil of bitter almonds contains a considerable proportion of Prussic Acid ; but the almonds themselves may be eaten with impunity, in moderation.

*Treatment.*

Ammonia, Brandy, and Oil of Turpentine in small doses, conjoined with artificial respiration if required.

---

## ANIMAL POISONS.

---

*SPANISH FLIES: (Lytta, Cantharidis.)*

*Symptoms.*

THEY very soon cause great pain and irritation about the urinary organs; difficulty and pain in making water, which is often reduced in quantity, though sometimes increased; breath has an unpleasant smell, and there is often a great aversion to liquids; stomach and bowels are tense and tender; satyriasis, sometimes in a very distressing degree; frequently bloody evacuations; blood is also frequently ejected by vomiting, and passed with the urine; sometimes convulsions and tetanus.

*Treatment.*

Diluent emetics ; mucilaginous draughts, in which Gum Arabic is dissolved, in abundance ; warm bath ; clysters of oil. The antiphlogistic treatment is generally required, with opiates.

*Morbid Appearances.*

Inflammation of the stomach and bowels ; also of the kidneys, ureters, and bladder ; sometimes the penis is in a gangrenous state.

*BITES OF SNAKES, VIPERS,  
VENOMOUS SERPENTS, &c.*

*Symptoms.*

Acute pain and swelling of the part bitten, which soon extends over the limb ; nausea and intoxicating symptoms soon come on, succeeded by delirium ; the part bitten becomes livid, and often gangrenous ; pulse quick and irregular ; breathing difficult and anxious ; often bilious vomiting ; sometimes impossibility of swallowing, with convulsions.

*Treatment.*

If possible remove the parts bitten, by excision ; then use some caustic application. Oil Ammonia are generally preferred : in the West Indies they employ Eau de Luce, the action of which depends on the Ammonia it contains. Give Brandy, Ammonia, and other stimuli, with Opiates. Arsenic has been recommended in doses of gr. ss. or gr. j., and it is said, that this practice has been successful. (Vide Medico-Chirurgical Transactions.)

The bites of some venomous insects sometimes occasion unpleasant symptoms ; but local applications will generally be sufficient to cure them. If possible, remove the poison ; oil applied to the part will often relieve ; at other times some evaporating spirituous lotion is necessary, to subdue the inflammation which exists.

*MUSCLES, LOBSTERS, CRABS  
AND OTHER FISH.*

*Symptoms.*

Uneasiness and pain about the stomach, with sickness and head-ache; vertigo; redness and swelling of the face; generally a species of nettle-rash all over the body; shortness of breath; rarely, cold extremities, delirium and convulsions.

*Treatment.*

Emetics, diluents, and purgatives; stimulants and opiates if necessary.

---

*Means to be resorted to in cases where animation is suspended, from drowning, hanging, or breathing deleterious airs.*

---

*FROM DROWNING.*

FIRST remove all clothes, and then convey the patient to a convenient and airy situation;

artificial respiration is now to be commenced, by inflating the lungs from the nose by a pair of bellows passed up one nostril; or with your own mouth, if no other be at hand: when the proper apparatus can be readily procured, that is best for the purpose; after each inflation the lungs must be again emptied, by pressure made on the chest. Wrap the body in warm blankets, and apply warmth to the body in any gradual manner. It is useless and improper to rub the body with any stimulating application. Apply hot water to the feet, or warm bricks. Introduce an elastic tube into the stomach, in order to convey stimulating fluids into that organ, as Brandy, Wine, &c.

When respiration becomes natural, we suspend our artificial operations; and as soon as the patient is able to swallow, give wine and water, and nourishing food. Never leave the person until he has perfectly recovered his senses. If Oxygen Gas be at hand, it may be employed. Electricity has been considered by some as a useful adjunct; it may be tried.

*FOR HANGING.*

A similar plan of treatment is necessary. Bleeding is often required here, from the jugular vein, to relieve the vessels of the brain and lungs; it should only be in small quantity.

*FROM NOXIOUS VAPORS.*

Similar treatment necessary. Here the temperature of the body is generally above the natural standard, and cold water should be suddenly dashed over the body, in addition to the other means. It is likewise very desirable, if possible, to substitute Oxygen Gas for the atmospheric air, in these cases.

When suffocation is occasioned by substances lodging in the air passage, and thus obstructing respiration, it is often necessary to perform the operation of Bronchotomy: here two methods have been recommended; one dividing the rings of the Trachea longitudinally; the other making an opening between the Thyroid and Cricoid Cartilages: each operation has its advocates, but I should prefer the former.









*Papaver  
Somniferum*

*Common  
White Poppy*

# I. PAPAVER SOMNIFERUM—PAPAVER ALBUM.

*White Poppy.*

(Capsulæ; et capsularum immaturarum succus concretus.

(Class XIII. POLYANDRIA.—Order I. MONOGYNIA.

*Natural Order.* RHÆADES.

(*Generic Character.* Corolla four-petalled. Calyx two-leaved. Capsule one-celled, opening by pores under the persistent stigma.

(*Specific Character.* Calyces and capsules smooth; leaves incised and embracing the stem.

THIS species of poppy is a native of Asia, and is found wild in the south of Europe, where the seeds had probably been accidentally scattered; it is also cultivated in this country, flowering in July.

The root is annual, tapering, and branched. The stalk is erect, three or four feet in height, branch-

ed, of a glaucous green color, round and cylindrical. The leaves are large, alternate, lobed, deeply cut into various segments, and embracing the stem. The flowers are large, terminal, and solitary; the calyx consists of two very smooth, ovate, concave segments, which fall when the flower expands: the petals are large, roundish, entire, somewhat undulated, and commonly of a white or purple color: the filaments are numerous, slender, shorter than the corolla, supporting erect, compressed anthers: the germen is roundish, with a many-rayed stigma: the capsule is smooth, large, and filled with a great many small seeds.

From cultivation and difference in soil, several varieties of the *Papaver Somniferum* are met with the double varieties are not at all inferior to the uncultivated plant.

Every part of the plant has the peculiar odor and taste of opium: but the milky juice, which is the active ingredient, resides chiefly in the capsules. The seeds however, when perfectly ripe, contain scarcely any of the narcotic principle, but

are chiefly composed of mucilage, and in their native soil are often used as an article of food ; they have a sweetish bland taste, somewhat like almonds.

## II. HYOSCYAMUS NIGER.

*Common Henbane.*

*Herba et Semina.*

*Class V. PENTANDRIA.—Order I. MONOGYNIA.*

*Natural Order. LURIDÆ.*

*Generic Character.* Corolla funnel-shaped, obtuse.  
*Stamina* inclined. Capsule covered with a lid,  
 two-celled.

*Specific Character.* Leaves embracing the stem,  
 sinuate ; flowers sessile.


This annual plant is a native of England, and grows by the road-sides and in waste uncultivated places, thriving best in rich soils. It flowers in June and July.

The root is long, compact, tapering, and fibrous. The stalk is about two feet in height, erect, cylindrical, woody and branched, beset with white hairs. The leaves are large, alternate, embracing the stem, downy, deeply sinuated, undulated,



*Hypocyanus*  
Viger

Black  
Herbaceous





and of a sea-green color. The flowers are simple, placed on terminal leafy spikes ; they consist of a short tube with an expanded limb, which is divided into five obtuse segments, of a straw color, and reticulated with purple veins: the calyx is tubular, permanent, and divided into five segments ; the filaments are tapering and downy at the base, supporting purple anthers, and are inserted into the tube of the corolla : the style is longer than the corolla, and ends in a blunt stigma : the capsule is globular, invested with the body of the calyx, and contains numerous irregular brown seeds.

The general appearance of this plant would almost lead us to suspect its deadly nature, and this is confirmed by its strong, disagreeable, and narcotic odor ; but it has scarcely any taste, and possesses but a slight degree of acrimony.

## III. CONIUM MACULATUM—CICUTA.

*Common Hemlock.*

Folia.

*Class V. PENTANDRIA.—Order II. DIGYNIA.**Natural Order. UMBELLATÆ.*

*Generic Character.* *Partial involucre* placed only on one side, three-leaved. *Fruit* nearly globular, five-streaked, notched on both sides.


*Specific Character.* Seeds striated.

HEMLOCK is a large biennial umbelliferous plant, growing in the neighbourhood of dunghills, ditches, and in moist shady places, flowering in June and July.

The root is fusiform, about as thick as the finger, yellowish externally and whitish within, exuding, when cut, a milky juice. The stalk is cylindrical, from three to five feet high, thick, hollow, branched, leafy, smooth, shining, and beset with purple spots. The lower leaves are



*Conium* | *Common*  
*Maculatum* | *Hemlock*



large, tripinnated, of a bright green color, standing upon long foot-stalks, which proceed from the joints of the stem ; the smaller or upper leaves are bipinnate. The flowers are forming open, numerous, umbels, which are both partial and universal. The involucre consists of from three to seven short, reflected, lance-shaped leaflets, white at the margin : the partial involucre is composed of three or four leaflets, which are disposed on the external side of the umbel. The flowers are small, composed of five petals, unequal, heart-shaped, and inclining inwards ; and an entire calyx ; stamina the length of the petals, supporting whitish anthers ; the styles are filiform, larger than the petals, diverging and terminating in round stigmas. The fruit is oval, striated, containing two brownish seeds.

## IV. CICUTA VIROSA.


*Water Hemlock.**Class V. PENTANDRIA.—Order II. DIGYNIA.**Natural Order. UMBELLATÆ.**Generic Character. Fruit subovate, sulcated.**Specific Character. Umbel opposite; petioles marginate, obtuse.*

ROOT perennial, thick, hollow, and beset with numerous fibres. Stalk thick, round, striated, smooth, and about four feet high. Leaves pinnated: leaflets usually in ternaries, spear-shaped and serrated. Flowers in large expanding umbels. Partial involucre composed of several short, bristle-shaped leaves. Calyx scarcely to be seen. Florets uniform, fertile; each consisting of five ovate, greenish white petals: filaments five, longer than the petals: anthers simple and purplish: stigmas simple; fruit egg-shaped.



*Cicuta*  
*virosa*

Water  
Hemlock









*Lactuca*  
*Tirosa*

*Strong's lettuce*  
*Lettuce*

## V. LACTUCA VIROSA.

*Strong-scented Lettuce.*

Herba.

*Class XIX. SYNGENESIA.—Order I. ÆQUALIS.**Natural Order. COMPOSITÆ SEMIFLOSCULOSÆ.*

*Generic Character.* Receptacle naked. Calyx imbricate, cylindrical, with a membranous margin. Pappus simple, stipitate. Seed even.

*Specific Character.* Leaves horizontal; carina pointed and toothed.

THIS plant grows on the banks of ditches, flowering in July and August.

The stalk is about three feet in height, erect, slender, round, prickly below, and smooth above. The leaves are smoothish and toothed, the lower ones obovate and undivided; those of the stalk smaller, often lobed, embracing the stem; middle rib having prickles on its under side. Bractes cordate and pointed. Flowers numerous, com-

pound, of a yellow color, furnished with small scaly leaves; calyx oblong, and composed of small lanceolate scales: the corolla consists of florets scarcely longer than the calyx. Seeds elliptical, compressed, black and striated.





*Atropa* | *Deadly*  
*Belladonna* | *Nightshade*

## VI. ATROPA BELLADONNA.

*Deadly Nightshade.*

Folia.

Class V. PENTANDRIA.—Order I. MONOGYNIA.

Natural Order. LURIDÆ.

*Generic Character.* Corolla bell-shaped. *Stamina* distant. *Berry* globular, two-celled.

*Specific Character.* Stalk herbaceous; leaves oval and entire.

THIS perennial plant is common in hot and temperate climates, especially in stony and shady situations. It grows in many parts of England, but is seldom to be met with in the neighbourhood of London. It flowers towards the latter end of June; its fruit ripens in September, and it is frequently cultivated in our gardens.

The root of the Belladonna is thick, long, and branched; from which proceed several herbaceous, cylindrical, branched stalks, from three to five feet in height, of a purplish color. The leaves

are in pairs, oval, entire, soft, pointed, and of different sizes. The flowers are pendant, supported on solitary and axillary peduncles: the calyx is monophyllous, deeply divided into five segments: corolla monopetalous, bell-shaped, partially divided into five lobes; it is of a dusky purplish color, and contains five stamens, whose filaments are inserted into the base of the corolla, supporting roundish anthers; a spheroidal germen supporting a style, with a divided stigma. The fruit is a roundish berry, contained within the calyx, of a blackish color and pulpy, having several kidney-shaped seeds.







*Aconitum*  
*Napellus*

Common  
Wolf's Bane

## VII. ACONITUM NAPELLUS.

*Wolf's Bane, Monkshood or Aconite.*

*Class XIII. POLYANDRIA.—Order III. TRIGYNIA.*

*Natural Order. MULTISILIQUEÆ.*

*Generic Character.* Calyx wanting. Petals five, the uppermost arched. Nectaries two, peduncled, recurved. Pods three or five.

*Specific Character.* Laciniæ of the leaves linear, broadest above, and gashed.

THIS perennial plant is a native of France, Germany, and Switzerland, growing in elevated situations : it is also frequently cultivated as an ornament to our gardens.

The root is fusiform. The stem several feet in height, erect and leafy. The lower leaves are lobed, and deeply cleft, placed on long petioles ; the upper ones are nearly sessile, of a dark green color above, and pale beneath. The flowers are placed on unifloral, axillary peduncles, and terminate the stem in a long spike ; the petals are of a deep violet color, the uppermost hooded, cover-

ing two curious nectaries ; the lateral ones roundish, and the lower elliptical : the filaments are spreading, supporting whitish anthers ; the germens are from three to five, with simple reflected stigmas.

The whole plant is very deleterious in its recent state ; it has a narcotic odor, and a pungent acrid taste, and the heat which it occasions in the mouth will continue for some minutes. The activity of the plant is much diminished by drying. The root is the most active part, but the leaves only are used medicinally.





*Datura* | *Common*  
*stramonium* | *Thorn Apple*

## VIII. DATURA STRAMMONIUM.

*Thorn Apple, James-town Weed.*

Herba.

*Class V. PENTANDRIA.—Order I. MONOGYNIA.*

*Natural Order. LURIDÆ.*

*Generic Character.* Corolla funnel-shaped, plaited.  
*Calyx* tubular, angular, deciduous. *Capsule*  
 four-valved.


*Specific Character.* Pericarps spinous, erect, oval ;  
 leaves ovate, glabrous.

THE Thorn Apple is an annual plant, and a native of America, but is now found growing in many places in the vicinity of London, and elsewhere, upon dunghills, and amongst the rubbish thrown from gardens, which generally contains some of its seeds; for it is frequently cultivated, and when it once takes possession of a soil, it is with difficulty extirpated: it flowers in July.

The stalk is thick, round, smooth, spreading, dichotomous above, and rising from two to three

feet in height. The leaves are of a dark green color, large, irregularly ovate, pointed at the extremity, angular, deeply indented, and supported by round foot-stalks. The flowers are large, white, axillary, solitary, placed on short erect peduncles: the calyx is one-leaved, tubular, pentangular, and four-toothed: the corolla is funnel-shaped, plaited, furnished with a long cylindrical tube, longer than the calyx. The filaments are slender, adhering to the tube, and support oblong flat anthers; the style is filiform, terminating in a short club-shaped stigma; the germen is oblong, and placed above the insertion of the corolla: the fruit is large, fleshy, ovate, beset with sharp spines, four-celled below, and two-celled above, containing numerous kidney-shaped seeds.







*Helleborus*  
*Niger*

*Black*  
*Hellebore*

## IX. HELLEBORUS NIGER.

*Black Hellebore or Christmas Rose.*

Radix.

*Class XIII. POLYANDRIA.—Order VI. POLYGYNIA.*

*Natural Order. MULTISILIQUEÆ.*

*Generic Character.* Calyx wanting. Petals five, or more. Nectaries bilabiate, tubular. Capsules many-seeded, nearly erect.

*Specific Character.* Scape one or two flowered, nearly naked; leaves pedate.

THIS plant is a native of Austria, the Apennines and Pyrenees, flowering from December to March. It is now cultivated in our gardens.

The root is perennial, transverse, rough, knotted, externally black, and internally whitish, sending off many depending fibres. The scapes, or flower stalks, are erect, round, towards the bottom reddish, and surrounded by an involucre. The leaves are of a deep green color, compound, of a peculiar shape, generally divided into five leaflets, and spring directly from the root, by long

footstalks; the leaflets are elliptical, smooth, coriaceous, and the upper half serrated: the floral leaves, which are oval and concave, supply the place of the calyx: the petals are large, roundish, concave and spreading, at first of a white color, with a tint of red, but by age they turn green. The nectaries are eight, tubular, bilabiate, and of a greenish color; filaments numerous, with yellow anthers; the germs vary from four to eight.

The roots of several plants have been occasionally mixed with those of the Black Hellebore, and sold as the genuine article; a fraud of the greatest importance to detect, as they possess properties widely different, and some of them are so very active that mischievous consequences have been the result of exhibiting them; for they cannot very readily be distinguished.

The fibrous part of the root, which alone is employed medicinally, is about the thickness of a straw, and six inches in length, of a deep brown color externally, and internally whitish; their taste is bitter and acrid, leaving a sensation of heat upon the tongue; their odor is nauseous and acrid, but much impaired by age.





*Helleborus*  
*viridifolius*

*Sinking*  
*Helleborus*

## X. HELLEBORUS FŒTIDUS.

*Fœtid Hellebore, or Bear's Foot.*

*Class XIII. POLYANDRIA.—Order VI. POLYGYNIA.*

*Natural Order. MULTISILIQUÆ.*

*Generic Character. Calyx none. Petals five or more. Nectaries bilabiate, tubular. Capsules many-seeded, nearly erect.*


*Specific Character. Stalk many-flowered, leafy; leaves pedate.*

THIS plant grows under hedges and in shady situations, flowering in March and April.

The root is small and bent, with numerous dark colored fibres. The stem is nearly two feet in height, round, strong, naked, and towards the top divided and subdivided into branches. The leaves are on long, channelled footstalks, surrounding the middle of the stem, and of a deep lurid green color, and pedate; the leaflets are long, narrow, lanceolate, serrate. At each ramification of the flower stem, are scaly, smooth, alternate, trifid

leaves ; those near the flowers are oval, entire and pointed. The flowers are numerous, terminal, peduncled and pendent : the petals are five oval, and concave, persistent, of a pale green color, the margins usually tinged with purple : stamina the length of the petals : anthers white : germens three, and resemble those of the *Helleborus Niger*.







*Primula*  
*Albom*

*White*  
*Hellebore*

## XI. VERATRUM ALBUM.

*White Hellebore.*

Radix.

*Class XXIII. POLYGAMIA.—Order I. MONŒCIA.**Natural Order. CORONARIÆ.*

*Generic Character.* Hermaphrodite. *Calyx* wanting. *Corolla* six-petalled. *Stamens* six. *Pistils* three. *Capsules* three, many-sided.

*Malc.* The same, rudiment of a pistil.

*Specific Character.* Stalk decomposed above; corollas erect.

THE root is perennial, fleshy, and fusiform, having strong fibres collected into a head. The stem is thick, round, hairy, erect and branching. The leaves are oblong, ovate, plaited longitudinally, of a yellowish green color, and embracing the stem at the base. The flowers are in long terminal spikes, composed of small alternate spikelets, each accompanied by a lanceolate bract: each flower consists of six persistent petals of a

pale green color : the filaments closely surround the germen, diverge, and terminate in yellow quadrangular anthers : the germens are three in each hermaphrodite flower, oblong, with erect bifid styles, crowned with flat spreading anthers.





*Colchicum*  
*Autumnale*



*Headed*  
*Saffron*

## XII. COLCHICUM AUTUMNALE.

*Common Meadow Saffron.*

Radix vel Bulbus Recens.

Class VI. HEXANDRIA.—Order III. TRIGYNIA.

*Natural Order. SPATHACEÆ.**Generic Character.* Corolla, six-parted, with a rooted tube. Capsules, connected, inflated.*Specific Character.* Leaves, flat, lanceolate, erect.

THIS perennial plant grows in moist meadow-grounds in the more temperate countries of Europe, flowering at the beginning of autumn without leaves, and bearing the fruit subsequently to the leaves.

The bulb is double, solid, succulent, and covered with a brown membranous coat. The leaves make their appearance in spring, and are radical, spear-shaped, and somewhat waved. The flower appears in autumn, immediately succeeding the decay of the leaves; it is large, of a purplish color, and springing from the root by a long naked

